



## REVIEW ARTICLE

### MANAGEMENT OF CHILDHOOD ASTHMA IN THE EMERGENCY DEPARTMENT IN MARRAKESH, MOROCCO

\*Soufiane EL MOUSSAOUI and Bourrous, M.

Child Emergency Department, Mother and Child Hospital, Mohammed VI University Hospital, Morocco

#### ARTICLE INFO

##### Article History:

Received 14<sup>th</sup> July, 2018  
Received in revised form  
17<sup>th</sup> August, 2018  
Accepted 10<sup>th</sup> September, 2018  
Published online 30<sup>th</sup> October, 2018

##### Keywords:

Asthma,  
Children,  
Emergency  
Management.

#### ABSTRACT

Asthma is the most common chronic disease in childhood. The morbidity and asthma mortality have become a problem in recent years, the most often due to inadequate therapeutic management. The work described in this study was directed at learning more about how asthma is managed in the community, and steps that can be taken to improve the management of these conditions. For this goal, we performed a prospective study with 216 patients; the population studied included children between the ages of 2 and 15 who presented to emergency department (ED) in The Mother-Child Hospital (Mohammed VI University hospital) in Marrakech over one year from January to December 2016. We found that. The majority of the patient (83.8%) had already developed an asthma attack. The personal and family history of asthma and /or atopy are present in more than half of the patients (57%). We also found that the allergens were the main precipitating factors (94%). The reason for consultation was cough in 94% of children. In Clinical examination, all patients had wheezes (100%) followed by Polypnea (63.4%). The treatment of choice was nebulization of salbutamol used for (96.3%) of children, followed by injection corticotherapy (56%) and oral corticotherapy (50%). Antibiotics were prescribed for 127 children (58.8%). In the result of this management 85.6% of patients presented an improvement and received ambulatory treatment. Childhood asthma remains under diagnosed and subcontracted it must be evoked as soon as the respiratory symptoms are repeated. In addition to the thoracic radiography which must be required systematically, the respiratory functional exploration not only contributes to the control of asthma but also for the evaluation of the effectiveness of the treatment prescribed, without forgets education during each consultation. Adhering to asthma regimes implies not only taking medications to relieve asthma attacks, but also adjusting their life styles in order to prevent asthma attacks.

Copyright © 2018, Soufiane EL MOUSSAOUI and Bourrous. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Asthma is the most common chronic disease in childhood, which affects >6.6 million children in the United States characterized by airway obstruction and hyper responsiveness. Asthma most often starts early in life and has variable courses and unstable phenotypes which may progress or remit over time (Molly Martin *et al.*, 2016). Morbidity and asthma mortality in children have become a problem in recent years (Aniba, 2011). The most often due to inadequate therapeutic management. Treatment of an asthma exacerbation is complex, involving a temporal and multi-disciplinary evaluation and reevaluation to adjust asthma medications and make a disposition decision. It is challenging to provide standardized care in a fast-paced and overcrowded environment like the ED. The evolution of asthma is relatively unknown. Indeed, 40%-50% of children followed for asthma will not be bothered by this disease as adults and nearly 50% of children follow for asthma will not disappear than disease after puberty (Aniba, 2011).

\*Corresponding author: Soufiane EL MOUSSAOUI,  
Child Emergency Department, Mother and Child Hospital, Mohammed VI University Hospital, Morocco.

The goal of this study is to assess the management of asthma and make an asthma care protocol in the pediatric ED to help standardize care and reduce time to disposition decision.

## PATIENTS AND METHODS

This work is a prospective study of children aged between 2 and 15 years, presented to the pediatric emergency department (ED) of University Hospital Mohammed VI Marrakech over one year from 1st January to 31 December 2016. This study included data collected through the exploitation records, filled out by duty doctors. We had simple and univariate data, processed by Microsoft Excel in duration of two weeks after consulting the epidemiological Laboratory of the Faculty of Medicine and Pharmacy Marrakech.

## RESULTS

The most frequent age of the asthma crisis in our study was between 2-6 year (59.25%) and an average age of 6.32 years, with a sex ratio of 1.5 with male predominance (60%). The

socioeconomic level was medium in (55%) of the cases and the residents were unhealthy in 23.1% of cases. More than half of the patients (57%) had a family history of asthma and/or atopy, most frequently in mothers (29%). In our study we found that the allergens were the main precipitating factors (94%) followed by physical exercise 81.9% than passive smoking 51.85%. The reason for consultation was: cough (104 children), dyspnea (84 children) or both (28 patients). C-reactive protein (CRP) and complete blood count (CBC) were requested respectively in 52 (24%) and 65 (30.1%). The chest X-ray was performed in 151 cases (69.9%) with the following results: normal in 24%, thoracic distension in 53% and infection focus in 43% of our patients. The treatment of choice was nebulization of salbutamol used for (96.3%) of children followed by injection Corticotherapy (56%) then oral Corticotherapy (50%). Antibiotics were prescribed for 58.8% of our children. The crisis severity was classified to mild in 80%, moderate in 18% and severe in 2% of the cases. We found also that 28.7% of children had regular follow of their asthma. The rest of the patients were distributed among poor compliance (26.8%), not followed (17%) and disregarded (27.5%). Finally we found that 185 children (85.6%) presented an improvement and received ambulatory treatment, 27 (12.5%) had been hospitalized on pediatric department A and only 4 (1.9%) was admitted in Intensive Care Unit (ICU).

**Table 1. Distribution of patients according to their personal history**

Personal History	Percentage
Followed for asthma	52%
Previous asthma attack	83.8%
Previous treatment	63.8%
Personal atopy	44%
Respiratory infection	65.27%
chronic vomiting	8.7%

**Table 2. Distribution of patients according to their functional signs**

Functional signs	Percentage
Dyspnea	97.9%
Dry cough	94%
Rhinorrhea	58.8%
Wheezing	30%
Sweating	12%
Cyanosis	3%
Disturbance of conscience	2%

**Table 3. Distribution of patients according to clinical signs**

Clinical signs	Percentage
polypnea	63.4%
Fever	32.9%
distended chest	38%
tachycardia	14.3%
subcutaneous emphysema	1.4%
decreased breath sounds	6%

**DISCUSSION**

The emergency department (ED) is where many patients with asthma seek and receive care. Respiratory disorders are the most common reason for ED visits in children after injuries and poisonings. Asthma is one of the most common chronic diseases among children in the United States, affecting more than 7 million children and costing more than 50 billion in direct health care costs annually (Molly Martin *et al.*, 2016). Not all children with asthma are affected equally. In 2010,

children with asthma in the United States experienced more than 900,000 asthma-related ED visits. Caring for asthma in EDs is costly for both families and the health care system. Some children who present to the ED for asthma care have severe asthma that is difficult to control under the best of circumstances, whereas other children have uncontrolled asthma due to improper management at home or in the ambulatory setting (Molly Martin *et al.*, 2016). In all situations, families require education and support while in the ED to safely and effectively transition their care to the ambulatory setting and home (Molly Martin *et al.*, 2016). With the shift from volume to value-based payment models for health care in the United States, there is increasing interest among health systems to identify and adopt strategies that improve the quality and outcomes of care transitions following ED discharge for asthma (Molly Martin *et al.*, 2016). Care transitions are defined as the movement of patients between health care practitioners, settings, and home as their need for health care evolves over time (Molly Martin *et al.*, 2016). The primary barriers to effective care transitions are inadequate communication, patient education, and accountability (Molly Martin *et al.*, 2016). ED providers frequently do not have access to ambulatory health records and cannot effectively coordinate management or arrange follow-up care. Patients and their caregivers often receive incomplete, confusing, or conflicting recommendations regarding care plans during health care transitions. This becomes even more challenging when patients lack a sufficient understanding of the disease or how to care for it. Accountability for the disease management falls either on the ambulatory provider or the patient/caregiver, both of whom often have incomplete information, limited resources to coordinate care, and may not communicate effectively with each other (Molly Martin *et al.*, 2016). Self-management education for asthma has been shown to improve patient outcomes and be a cost-effective component of asthma care in various settings (Molly Martin *et al.*, 2016).

However, studies of asthma self-management education interventions in the ED are limited. In an Agency for Healthcare Research and Quality review of self-management interventions for pediatric asthma through 2006, none of the 75 interventions identified in the review that directly targeted self-management and patient education were conducted in the ED (Molly Martin *et al.*, 2016). A Cochrane review of educational interventions for pediatric asthma in the ED included 38 studies but only 4 of these interventions actually occurred in the ED (Molly Martin *et al.*, 2016). Although the Cochrane review reported that educational intervention was strongly associated with a reduced risk for the subsequent ED visits (Molly Martin *et al.*, 2016). Only 1 of the 4 studies that delivered the educational intervention in the ED setting reported some improvements in health care utilization (Molly Martin *et al.*, 2016). The prevalence of asthma increased by approximately 5% annually in recent years. In December 2009 Asthma Insights and Reality in Maghreb (AIRMAG) found that the prevalence of children with asthma ranged from 3.5% in Tunisia, 4.4% in Morocco, and 6.4% in Algeria. The prevalence of asthma in the region of Safi was 3.4% in 2010 (Aniba, 2011). Numerous potential risk factors have been studied in relation to the development of asthma. Atopy is frequently identified as a strong risk factor for the development of asthma (Busse *et al.*, 2011). Our study objectified that 44% had a personal or family history of atopy. Martinez and al studies in USA found the influence of several genes including one major gene transmitted respectively by an autosomal co-

dominant and autosomal recessive (Piacentini *et al.*, 2014). In our study, the family history of asthma and/or atopy was recorded in more than half of cases, most frequently in mothers (29%). Air pollution and viral infections are well-established triggers for asthma exacerbations (Patel and Miller *et al.*, 2009). Recent studies have found that over 80% of wheezing episodes were associated with viral respiratory infections (Mackenzie *et al.*, 2013). In more than 60% of these children, the respiratory syncytial virus (RSV) was detected. The close link between bronchiolitis induced by viruses and the development of asthma has been demonstrated in several studies (Murray *et al.*, 1992). In our study, more than half of the children (57%) followed for asthma had a history of recurrent respiratory viral infections. The study of Platts-Mills and al showed, in patients allergic to dust mites, improved clinical signs and decreased bronchial hyper responsiveness (BHR), after a sojourn of two months in a hospital without mites (Platts mills *et al.*, 1999). In our study, we found that the allergens were the main precipitation factors, present in 94% of our patients. Parental smoking is a significant risk factor for acute lower respiratory tract infections in infants, and the development of wheezing and asthma in children. In our population, 51.8% of asthmatic children were exposed to parental smoking (Kanoh *et al.*, 2012). The classic triad of asthma includes cough, wheeze, and dyspnea. However, patients often present with only 1 of these symptoms which can make diagnosis challenging.

In our study, Dyspnea was found in 97.9% and dry cough in 94% of children. Pediatric status asthmaticus (PSA) is a medical emergency warranting prompt recognition and intervention. A status asthmaticus or severe asthma exacerbation is defined as an acute episode that does not respond to standard treatment with short acting  $\beta_2$ -agonists and corticosteroids. In a USA cohort, admission for status asthmaticus between 1992 and 2006 approximately halved from 1.92 to 0.93 per 1000 children. In contrast, ICU care related to asthma increased from 0.09 to 0.31 per 1000 patients. In our study, 4 patients presented a status asthmaticus and were admitted to ICU. The clinical expression of Exercise induced asthma (EIA) is that of asthma attack whose only characteristic is to follow an exercise. Typically, it occurs in the exercise, in a subject with normal respiratory function, and reaches its maximum intensity 5 to 10 minutes after cessation of exercise (De Blic *et al.*, 2005). In our study, the physical exercise induced asthma attacks was found in 81.9% of children. Although, diagnosis of asthma is clinic, chest X-ray is requested systematically to eliminate false asthma. The chest X-ray was performed in 69.9% in our children. The respiratory function test is the examination that enables both better diagnostic approach to asthma and therapeutic monitoring. The use of peak respiratory flow has been widely adopted for monitoring patients with asthma. Reduced peak flow measurements do not differentiate between obstructive and restrictive diseases, spirometry and sometimes measurement of lung volumes are necessary to distinguish the two (Quanjer and Lebowitz, 1997). In our study, both of them (FEV and PEF) are not yet available in the emergency department. Treatment of asthma should be started immediately. The goals of therapy include correction of hypoxemia, rapid reversal of airway obstruction, and treatment of inflammation. Short-acting  $\beta$ -agonists are the recommended first-line therapy for the acute asthma exacerbation. A meta-analysis including 21 pediatric studies and more than 2000 children demonstrated that in acute asthma spacers were as effective as nebulizers in

limiting hospitalization rates and reducing the time spent in emergency department (Plotnick and Ducharme, 2008). In our study, the majority of children (96%) used nebulized salbutamol. Corticosteroids are recommended for asthma exacerbations that are incompletely responsive to inhaled  $\beta$ -agonists. In our study, the use of oral corticosteroids was found in the half of children (50.5%). For antibiotics, The recent National Asthma Education and Prevention Program Expert Panel Report indicates that antibiotics are not currently recommended for the treatment of acute asthma exacerbations except when fever, purulent sputum or clear evidence of infection are present. We saw in our study a significant use of the antibiotic in the treatment of asthma in more than half (58.8%) of the cases. In addition to these treatments, the magnesium sulfate is a potential therapeutic agent in asthma because of its bronchodilating effect on smooth muscle cells and reduction of the neutrophilic burst associated with inflammation. In our emergency department, the magnesium sulfate was not used in any of our patients. Asthma control refers to control of the clinical manifestations of the disease, and it is the ultimate goal of asthma management. There is a clear relationship between asthma severity and asthma control (Humbert *et al.*, 2007). In our study, we found that 62 children (28.7%) had regular control of their asthma. The long-term prognosis of asthma remains relatively unknown as there are few prospective studies from childhood to adulthood. The Blair Study of 200 asthmatic children showed that two-thirds of children with mild asthma had an excellent state twenty years later: life is normal, their attacks are absent or rare and very sensitive to bronchodilators (Roorda *et al.*, 1993). In contrast, two-thirds of severe asthma remain so 20 years later.

## Conclusion

In light of our results, it is clear that there was no specific protocol for the management of asthma attack in our pediatric ED. We found that there were certain key aspects of the management of childhood asthma that were being overlooked. We speculated that a local asthma educational programme intended to reduce the number of ED visits for asthma exacerbations. Further prospective studies are needed to corroborate this hypothesis.

## REFERENCES

- Aniba J. 2011. Prévalence de l'asthme chez l'enfant dans la région de Safi. Thèse Doctorat Médecine, Marrakech, n° 41, 53 pages.
- Busse WW, *et al.* 2011. Expert Panel Report Three. Guidelines for the diagnosis and management of asthma. *National Institutes of Health*, 12:160.17.
- Care transition interventions for children with asthma in the emergency department Molly A. Martin, MD *et al.* JACI Journal Club 2016.
- De Blic J. 2005. Asthma in children and infants. *Rev Mal Resp*, Av; 22(2):19-2.
- Humbert M, Holgate S, Boulet LP, *et al.* 2007 Asthma control or severity: that is the question. *Allergy*, 62(2): 95-101.
- Kanoh, M., Kaneita, Y., Hara, M., Harada, S., Gon, Y., Kanamaru, H., Ohida, 2012. Parental smoking habits and development of asthma in early childhood. *Prev. Med.*, 54, 94-96.
- Mackenzie KJ, Anderton SM, Schwarze J. 2013. Viral respiratory tract infections and asthma in early life; cause and effect?. *Clin Exp Allergy*, 44(1):9-19.

- Murray M, Webb MSC, O'callagan C *et al.* 1992. Respiratory status and allergy after bronchiolitis. *Arch Dis Child.*, 67:482.
- Patel MM, Miller RL. 2009. Air pollution and childhood asthma: recent advances and future directions. *Curr Opin Pediatr.*, 21:235-42.
- Piacentini S, A Verrotti, Polimanti R, *et al.* 2014. Polymorphismes fonctionnels des gènes GSTA1 et GSTO2 associés à l'asthme chez les enfants asthmatiques en Italie. *Clin Exp Pharmacol Physiol.*, 41(3):180-4.
- Platts mills TA., Tovey ER, Mitchell EB, *et al.* 1999. Reduction of bronchial hyperreactivity during prolonged allergen avoidance. *Lancet.*, 2:675-8.
- Plotnick L, Ducharme F. 2008. Combined inhaled anticholinergics and beta2-agonists for initial treatment of acute asthma in children. *Cochrane Database Syst Rev.*
- Quanjer P.H., M.D. Lebowitz, 1997. Peak expiratory flow: conclusions and recommendations of a Working Party of the European Respiratory Society. *Eur Respir J.*, 10(24):2-8.
- Roorda RJ, Gerritsen J, Vanaaldren, *et al.* 1993. Risk factors for persistence of respiratory symptoms in childhood asthma. *Am Rev Respir Dis.*, 148:1490-1495.

\*\*\*\*\*